

**Author:** Michael Jibson, M.D., Ph.D., 2009

License: Unless otherwise noted, this material is made available under the terms of the **Creative Commons Attribution–Share Alike 3.0 License:**  
<http://creativecommons.org/licenses/by-sa/3.0/>

**We have reviewed this material** in accordance with U.S. Copyright Law **and have tried to maximize your ability to use, share, and adapt it.** The citation key on the following slide provides information about how you may share and adapt this material.

Copyright holders of content included in this material should contact [open.michigan@umich.edu](mailto:open.michigan@umich.edu) with any questions, corrections, or clarification regarding the use of content.

For more information about **how to cite** these materials visit <http://open.umich.edu/education/about/terms-of-use>.

Any **medical information** in this material is intended to inform and educate and **is not a tool for self-diagnosis** or a replacement for medical evaluation, advice, diagnosis or treatment by a healthcare professional. Please speak to your physician if you have questions about your medical condition.

**Viewer discretion is advised:** Some medical content is graphic and may not be suitable for all viewers.

# Citation Key

for more information see: <http://open.umich.edu/wiki/CitationPolicy>

## Use + Share + Adapt

{ Content the copyright holder, author, or law permits you to use, share and adapt. }



**Public Domain – Government:** Works that are produced by the U.S. Government. (17 USC § 105)



**Public Domain – Expired:** Works that are no longer protected due to an expired copyright term.



**Public Domain – Self Dedicated:** Works that a copyright holder has dedicated to the public domain.



**Creative Commons – Zero Waiver**



**Creative Commons – Attribution License**



**Creative Commons – Attribution Share Alike License**



**Creative Commons – Attribution Noncommercial License**



**Creative Commons – Attribution Noncommercial Share Alike License**



**GNU – Free Documentation License**

## Make Your Own Assessment

{ Content Open.Michigan believes can be used, shared, and adapted because it is ineligible for copyright. }



**Public Domain – Ineligible:** Works that are ineligible for copyright protection in the U.S. (17 USC § 102(b)) \*laws in your jurisdiction may differ

{ Content Open.Michigan has used under a Fair Use determination. }



**Fair Use:** Use of works that is determined to be Fair consistent with the U.S. Copyright Act. (17 USC § 107) \*laws in your jurisdiction may differ

Our determination **DOES NOT** mean that all uses of this 3rd-party content are Fair Uses and we **DO NOT** guarantee that your use of the content is Fair.

To use this content you should **do your own independent analysis** to determine whether or not your use will be Fair.

# Personality and Personality Disorders

## M2 Psychiatry Sequence

Michael Jibson  
Fall 2008



# Personality Traits

---

- “Enduring patterns of perceiving, relating to, and thinking about the environment and oneself that are exhibited in a wide range of social and personal contexts.” (DSM-IV)

# Personality Disorder

---

- “Only when personality traits are inflexible and maladaptive and cause significant functional impairment or subjective distress do they constitute Personality Disorders.” (DSM-IV)

# Diagnostic Criteria for a Personality Disorder

## General Diagnostic Criteria for a Personality Disorder (DSM-IV)

- A. An enduring pattern of inner experience and behavior that deviates markedly from the expectations of the individual's culture. This pattern is manifested in two (or more) of the following areas:
- (1) cognition (i.e., ways of perceiving and interpreting self, other people and events)
  - (2) affectivity (i.e., the range, intensity, lability, and appropriateness of emotional response)
  - (3) interpersonal functioning
  - (4) impulse control
- B. The enduring pattern is inflexible and pervasive across a broad range of personal and social situations.
- C. The enduring pattern leads to clinically significant distress or impairment in social, occupational, or other important areas of functioning.
- D. The pattern is stable and of long duration and its onset can be traced back at least to adolescence or early adulthood.
- E. The enduring pattern is not better accounted for as a manifestation or consequence of another mental disorder.
- F. The enduring pattern is not due to the direct physiological effects of a substance (e.g., a drug of abuse, a medication) or a general medical condition (e.g., head trauma).

# Dimensions of Personality

## Hippocrates - “Four humors”

---

- Blood - Emotional lability
  - Black bile - Depression
  - Yellow bile - Anger
  - Phlegm - Slow, stolid, cold
-

# Dimensions of Personality

## Carl Jung - *Psychological Types* (1921)

---

- Introvert-Extravert
  - Thinking-Feeling
  - Sensing-Intuiting
  - Judging-Perceiving
-



# Dimensions of Personality

## Assessment Instruments

---

- Self-report inventories
    - Minnesota Multiphasic Personality Inventory (MMPI, 1937)
      - >500 True-false questions (“I believe I am being plotted against.” “I sometimes tease animals.”)
      - Ten clinical scales (hypochondriasis, depression,...)
      - Three validity scales (detect “faking good” and “faking bad”)
      - Four special scales (ego strength, anxiety,...)
-

# Dimensions of Personality

## Assessment Instruments

---

- Self-report inventories
    - Meyers-Briggs - Scores are plotted along Jung's four dimensions
-

# Dimensions of Personality

## Assessment Instruments

---

- Structured clinical interview for diagnosis (SCID) - Based on diagnostic criteria, not dimensions
  - Clinical interview
-

# Dimensions of Personality

## Assessment Instruments

---

- Projective tests - Not diagnostic, but show patterns of thought, dynamics, defenses, disorders of thought, etc.
    - Rorschach (ink-blot)
    - Thematic Apperception Test (TAT) - (tell stories about evocative pictures)
    - Sentence-Completion Test (SCT) - (“I like...” “Sometimes I wish...”)
    - Draw-A-Person (DAP)
-

# Plate I of the Rorschach Test



# Etiology

## Genetic and biologic factors

---

- Concordance rates of personality traits for monozygotic twins are higher than for dizygotic twins, even if they are raised apart
-

# Etiology - Genetic and Biologic Factors

## Larry Siever

---

- Cognitive disorganization (includes “interpersonal detachment”) - Cluster A
  - Impulsivity - Cluster B
    - Decreased 5-HT and 5-HIAA (5-HT metabolite)
  - Affective instability - Cluster B
    - Hyperresponsivity of noradrenergic system
  - Anxiety / Inhibition - Cluster C
    - High autonomic arousal from infancy
-

# Etiology - Genetic and Biologic Factors

## Robert Cloninger

---

- Novelty seeking
  - Harm avoidance
  - Reward dependence
-



# Etiology

## Environmental factors

---

- Parenting and family style
  - Psychosocial milieu
-

# Etiology

## Psychodynamic factors

---

- Internal drives and defenses
  - Developmental tasks and stages
-

# Cluster A Personality Disorders

- Odd or Eccentric

# Cluster A Personality Disorders

## Paranoid Personality Disorder

---

- “A pattern of distrust or suspiciousness such that others’ motives are interpreted as malevolent.” (DSM-IV)
-

# Cluster A Personality Disorders

## Diagnostic Criteria for Paranoid Personality Disorder (DSM-IV)

- A. A pervasive distrust and suspiciousness of others such that their motives are interpreted as malevolent, beginning by early adulthood and present in a variety of contexts, as indicated by four (or more) of the following:
- (1) suspects, without sufficient basis, that others are exploiting, harming, or deceiving him or her
  - (2) is preoccupied with unjustified doubts about the loyalty or trustworthiness of friends or associates
  - (3) is reluctant to confide in others because of unwarranted fear that the information will be used maliciously against him or her
  - (4) reads hidden demeaning or threatening meanings into benign remarks or events
  - (5) persistently bears grudges, i.e., is unforgiving of insults, injuries, or slights
  - (6) perceives attacks on his or her character or reputation that are not apparent to others and is quick to react angrily or to counterattack
  - (7) has recurrent suspicions, without justification, regarding fidelity of spouse or sexual partner
- B. Does not occur exclusively during the course of Schizophrenia, a Mood Disorder With Psychotic Features, or another Psychotic Disorder and is not due to the direct physiological effects of a general medical condition.

# Cluster A Personality Disorders

## Paranoid Personality Disorder

---

- Prevalence: 2% of population.
  - Sex ratio: F:M=3:1
  - Comorbidity: Brief reactive psychosis, delusional disorder, anxiety, substance abuse, depression, schizophrenia
  - Family: Delusional disorder, schizophrenia, Cluster A disorders
-

# Cluster A Personality Disorders

## Paranoid Personality Disorder

---

- Treatment
    - Psychotherapy - Treatment of choice, but patients have limited introspection
    - Medication - Anxiolytics are often useful; antipsychotics sometimes helpful
-

# Cluster A Personality Disorders

## Paranoid Personality Disorder

---

- Physician-patient interaction
    - A straightforward approach, without an expectation of personal warmth is most effective
    - Greater empathy may actually make the patient more anxious
-



# Cluster A Personality Disorders

## Schizoid Personality Disorder

---

- “A pattern of detachment from social relationships and a restricted range of emotional expression.” (DSM-IV)
-

# Cluster A Personality Disorders

## Diagnostic Criteria for Schizoid Personality Disorder (DSM-IV)

- A. A pervasive pattern of detachment from social relationships and a restricted range of expression of emotions in interpersonal settings, beginning by early adulthood and present in a variety of contexts, as indicated by four (or more) of the following:
- (1) neither desires nor enjoys close relationships, including being part of a family
  - (2) almost always chooses solitary activities
  - (3) has little, if any, interest in having sexual experiences with another person
  - (4) takes pleasure in few, if any, activities
  - (5) lacks close friends or confidants other than first-degree relatives
  - (6) appears indifferent to the praise or criticism of others
  - (7) shows emotional coldness, detachment, or flattened affectivity
- B. Does not occur exclusively during the course of Schizophrenia, a Mood Disorder With Psychotic Features, another Psychotic Disorder, or a Pervasive Development Disorder and is not due to the direct physiological effects of a general medical condition.

# Cluster A Personality Disorders

## Schizoid Personality Disorder

---

- Prevalence: 3% of population
  - Sex ratio: M>F
  - Comorbidity: Delusional disorder, schizophrenia
  - Family: Schizophrenia, Cluster A disorders, esp. schizotypal personality disorder
-

# Cluster A Personality Disorders

## Schizoid Personality Disorder

---

- Treatment

- Psychotherapy - Treatment of choice; introspection is usually good
  - Medication - Low doses of antipsychotics or antidepressants are occasionally helpful
-

# Cluster A Personality Disorders

## Schizoid Personality Disorder

---

- Physician-patient interaction
    - A straightforward approach, without an expectation of personal warmth is preferred
    - Greater empathy may actually make the patient more anxious
-

# Cluster A Personality Disorders

## Schizotypal Personality Disorder

---

- “A pattern of acute discomfort in close relationships, cognitive or perceptual distortions, and eccentricities of behavior.” (DSM-IV)
-

# Cluster A Personality Disorders

## Diagnostic Criteria for Schizotypal Personality Disorder (DSM-IV)

- A. A pervasive pattern of social and interpersonal deficits marked by acute discomfort with, and reduced capacity for, close relationships as well as by cognitive or perceptual distortions and eccentricities of behavior, beginning by early adulthood and present in a variety of contexts, as indicated by five (or more) of the following:
- (1) ideas of reference (excluding delusions of reference)
  - (2) odd beliefs or magical thinking that influences behavior and is inconsistent with subcultural norms (e.g., superstitiousness, belief in clairvoyance, telepathy, or "sixth sense"; in children and adolescents, bizarre fantasies or preoccupations)
  - (3) unusual perceptual experiences, including bodily illusions
  - (4) odd thinking and speech (e.g., vague, circumstantial, metaphorical, overelaborate, or stereotyped)
  - (5) suspiciousness or paranoid ideation
  - (6) inappropriate or constricted affect
  - (7) behavior or appearance that is odd, eccentric, or peculiar
  - (8) lack of close friends or confidants other than first-degree relatives
  - (9) excessive social anxiety that does not diminish with familiarity and tends to be associated with paranoid fears rather than negative judgments about self

# Cluster A Personality Disorders

## Schizotypal Personality Disorder

---

- Prevalence: 3% of population, but uncommon in clinical settings.
  - Sex ratio: M>F
  - Comorbidity: Depression, anxiety, brief reactive psychosis, delusional disorder, schizophrenia
  - Family: Schizophrenia, Cluster A disorders.
-



# Cluster A Personality Disorders

## Schizotypal Personality Disorder

---

- Treatment

- Psychotherapy - Treatment of choice. Insight may be limited.
  - Medication - Antipsychotics may be useful
-

# Cluster A Personality Disorders

## Schizotypal Personality Disorder

---

- Physician-patient interaction
    - A straightforward approach, without an expectation of personal warmth is preferred
    - Greater empathy may actually make the patient more anxious
    - Care must be taken not to ridicule odd or over-valued ideas
    - Avoid overt rejection - even a limited personal interaction may be very important to the patient, and its loss distressing
-

# Cluster B Personality Disorders

- Dramatic, Emotional, or Erratic

# Cluster B Personality Disorders

## Antisocial Personality Disorder

---

- “A pattern of disregard for, and violation of, the rights of others.” (DSM-IV)
-

# Cluster B Personality Disorders

## Diagnostic Criteria for Antisocial Personality Disorder (DSM-IV)

- A. There is a pervasive pattern of disregard for and violation of the rights of others occurring since age 15 years, as indicated by three (or more) of the following:
- (1) failure to conform to social norms with respect to lawful behaviors as indicated by repeatedly performing acts that are grounds for arrest
  - (2) deceitfulness, as indicated by repeated lying, use of aliases, or conning others for personal profit or pleasure
  - (3) impulsivity or failure to plan ahead
  - (4) irritability and aggressiveness, as indicated by repeated physical fights or assaults
  - (5) reckless disregard for safety of self or others
  - (6) consistent irresponsibility, as indicated by repeated failure to sustain consistent work behavior or honor financial obligations
  - (7) lack of remorse, as indicated by being indifferent to or rationalizing having hurt, mistreated, or stolen from another
- B. The individual is at least age 18 years.
- C. There is evidence of Conduct Disorder with onset before age 15 years.
- D. The occurrence of antisocial behavior is not exclusively during the course of Schizophrenia or a Manic Episode.

# Cluster B Personality Disorders

## Antisocial Personality Disorder

---

- Prevalence: 3% of males and 1% of females
  - Sex ratio: M:F=3:1
  - Comorbidity: Substance abuse, attention deficit disorder, depression, anxiety
  - Family: Somatization disorder, substance abuse, Cluster B disorders, esp. antisocial personality disorder
-

# Cluster B Personality Disorders

## Antisocial Personality Disorder

---

- Major clinical issues
    - Violence
    - Criminal behavior
    - Suicide
-

# Cluster B Personality Disorders

## Antisocial Personality Disorder

---

- Treatment – no psychiatric treatment addresses the core pathology
    - Psychotherapy - Not generally useful, although it may alleviate depression and anxiety, especially if the patient is immobilized (e.g., in jail)
    - Medication - May be useful for comorbid disorders; uncontrolled rage may be helped somewhat by antipsychotics or mood stabilizers
-



# Cluster B Personality Disorders

## Antisocial Personality Disorder

---

- Physician-patient interaction
    - Firm limits are essential
    - Substance abuse is a major problem
    - complicated by genuine distress and incessant manipulation
-

# Cluster B Personality Disorders

## Borderline Personality Disorder

---

- “A pattern of instability in interpersonal relationships, self-image, and affects, and marked impulsivity.” (DSM-IV)
-

# Cluster B Personality Disorders

## Diagnostic Criteria for Borderline Personality Disorder (DSM-IV)

A pervasive pattern of instability of interpersonal relationships, self-image, and affects, and marked impulsivity beginning by early adulthood and present in a variety of contexts, as indicated by five (or more) of the following:

- (1) frantic efforts to avoid real or imagined abandonment
- (2) a pattern of unstable and intense interpersonal relationships characterized by alternating between extremes of idealization and devaluation
- (3) identity disturbance: markedly and persistently unstable self-image or sense of self
- (4) impulsivity in at least two areas that are potentially self-damaging (e.g., spending, sex, substance abuse, reckless driving, binge eating)
- (5) recurrent suicidal behavior, gestures, or threats, or self-mutilating behavior
- (6) affective instability due to a marked reactivity of mood (e.g., intense episodic dysphoria, irritability, or anxiety usually lasting a few hours and only rarely more than a few days)
- (7) chronic feelings of emptiness
- (8) inappropriate, intense anger or difficulty controlling anger (e.g., frequent displays of temper, constant anger, recurrent physical fights)
- (9) transient, stress-related paranoid ideation or severe dissociative symptoms

# Cluster B Personality Disorders

## Borderline Personality Disorder

---

- Prevalence: 3% of females and 1% of males
  - Sex ratio: F:M=3:1
  - Comorbidity: Depression, substance abuse, eating disorders, brief reactive psychosis
  - Family: Mood disorders, substance abuse, Cluster B disorders, esp. antisocial personality disorder
-

# Cluster B Personality Disorders

## Borderline Personality Disorder

---

- Major clinical issues
    - Suicide and self-mutilation
    - Splitting - seeing the world as all good or all bad
    - Rage
    - Psychosis
    - Childhood trauma (especially sexual)
    - Dissociation - depersonalization, derealization, amnestic episodes,
-

# Cluster B Personality Disorders

## Borderline Personality Disorder

---

- Treatment
    - Psychotherapy - Dialectical/behavioral therapy (DBT) is preferred. Individual, group, and cognitive/behavioral therapy (CBT) are difficult, but may be useful.
    - Medication - Low-dose antipsychotics, mood stabilizers, and standard-dose antidepressants are moderately useful. Anxiolytics are beneficial in a minority of patients.
-

# Cluster B Personality Disorders

## Borderline Personality Disorder

---

- Physician-patient interaction
    - Idealization, devaluation, and splitting are common
    - Firm limits and high tolerance for regressive (childish) behavior are essential
    - Countertransference must be monitored carefully
-

# Cluster B Personality Disorders

## Histrionic Personality Disorder

---

- “A pattern of excessive emotionality and attention seeking.” (DSM-IV)
-



# Cluster B Personality Disorders

## Diagnostic Criteria for Histrionic Personality Disorder (DSM-IV)

A pervasive pattern of excessive emotionality and attention seeking, beginning by early adulthood and present in a variety of contexts, as indicated by five (or more) of the following:

- (1) is uncomfortable in situations in which he or she is not the center of attention
- (2) interaction with others is often characterized by inappropriate sexually seductive or provocative behavior
- (3) displays rapidly shifting and shallow expression of emotions
- (4) consistently uses physical appearance to draw attention to self
- (5) has a style of speech that is excessively impressionistic and lacking in detail
- (6) shows self-dramatization, theatricality, and exaggerated expression of emotion
- (7) is suggestible, i.e., easily influenced by others or circumstances
- (8) considers relationships to be more intimate than they actually are

# Cluster B Personality Disorders

## Histrionic Personality Disorder

---

- Prevalence: 2-3% of population
  - Sex ratio: F>M
  - Comorbidity: Somatization and conversion disorders, depression, anxiety
  - Family: Cluster B disorders
-

# Cluster B Personality Disorders

## Histrionic Personality Disorder

---

- Treatment
    - Psychotherapy - Dynamic therapy is the treatment of choice
    - Medication - Antidepressants and anxiolytics may help comorbid depression and anxiety
-

# Cluster B Personality Disorders

## Histrionic Personality Disorder

---

- Physician-patient interaction
    - Overly dependent or seductive behavior is common
    - A dramatic presentation may obscure the differences between major and minor physical problems
    - Rapid fluctuation between overwhelming anxiety about a medical problem and total indifference is common.
-

# Cluster B Personality Disorders

## Narcissistic Personality Disorder

---

- “A pattern of grandiosity, need for admiration, and lack of empathy.” (DSM-IV)
-

# Cluster B Personality Disorders

## Diagnostic Criteria for Narcissistic Personality Disorder (DSM-IV)

A pervasive pattern of grandiosity (in fantasy or behavior), need for admiration, and lack of empathy, beginning by early adulthood and present in a variety of contexts, as indicated by five (or more) of the following:

- (1) has a grandiose sense of self-importance (e.g., exaggerates achievements and talents, expects to be recognized as superior without commensurate achievements)
- (2) is preoccupied with fantasies of unlimited success, power, brilliance, beauty, or ideal love
- (3) believes that he or she is "special" and unique and can only be understood by, or should associate with, other special or high-status people (or institutions)
- (4) requires excessive admiration
- (5) has a sense of entitlement, i.e., unreasonable expectations of especially favorable treatment or automatic compliance with his or her expectations
- (6) is interpersonally exploitative, i.e., takes advantage of others to achieve his or her own ends
- (7) lacks empathy: is unwilling to recognize or identify with the feelings and needs of others
- (8) is often envious of others or believes that others are envious of him or her
- (9) shows arrogant, haughty behaviors or attitudes

# Cluster B Personality Disorders

## Narcissistic Personality Disorder

---

- Prevalence: <1% of population
  - Sex ratio: 50-75% male
  - Comorbidity: Mood disorders, anorexia, substance abuse
  - Family: Cluster B disorders
-

# Cluster B Personality Disorders

## Narcissistic Personality Disorder

---

- Treatment
    - Psychotherapy - Psychotherapy is difficult, but is the treatment of choice. Interpersonal therapy (IPT) may be most effective.
    - Medication - Antidepressants or mood stabilizers may be helpful for comorbid mood disorders
-



# Cluster B Personality Disorders

## Narcissistic Personality Disorder

---

- Physician-patient interaction
    - Idealization gives way rapidly to contemptuous devaluation
    - Entitlement and condescension are common
    - Be aware that it is more often the physician than the patient who has these traits
-

# Cluster B Personality Disorders

- Anxious or Fearful

# Cluster C Personality Disorders

## Avoidant Personality Disorder

---

- “A pattern of social inhibition, feelings of inadequacy, and hypersensitivity to negative evaluation.” (DSM-IV)
-

# Cluster C Personality Disorders

## Diagnostic Criteria for Avoidant Personality Disorder (DSM-IV)

A pervasive pattern of social inhibition, feelings of inadequacy, and hypersensitivity to negative evaluation, beginning by early adulthood and present in a variety of contexts, as indicated by four (or more) of the following:

- (1) avoids occupational activities that involve significant interpersonal contact, because of fears of criticism, disapproval, or rejection
- (2) is unwilling to get involved with people unless certain of being liked
- (3) shows restraint within intimate relationships because of the fear of being shamed or ridiculed
- (4) is preoccupied with being criticized or rejected in social situation
- (5) is inhibited in new interpersonal situations because of feelings of inadequacy
- (6) views self as socially inept, personally unappealing, or inferior to others
- (7) is unusually reluctant to take personal risks or to engage in any new activities because they may prove embarrassing

# Cluster C Personality Disorders

## Avoidant Personality Disorder

---

- Prevalence: 0.5-1% of population
  - Sex ratio: M=F
  - Comorbidity: Social phobia, depression, anxiety
  - Family: Cluster C disorders
-

# Cluster C Personality Disorders

## Avoidant Personality Disorder

---

- Treatment
    - Psychotherapy - Individual and group therapy, typically interpersonal (IPT), is the treatment of choice
    - Medication - Antidepressants and anxiolytics are often useful for accompanying depression and anxiety
-

# Cluster C Personality Disorders

## Avoidant Personality Disorder

---

- Physician-patient interaction
    - Unconditional respect and concern are very helpful
    - Avoid implications of rejection
    - Be aware that even a limited personal interaction may be very important, and its loss very distressing
-

# Cluster C Personality Disorders

## Dependent Personality Disorder

---

- “A pattern of submissive and clinging behavior related to an excessive need to be taken care of.” (DSM-IV)
-



# Cluster C Personality Disorders

## Diagnostic Criteria for Dependent Personality Disorder (DSM-IV)

A pervasive and excessive need to be taken care of that leads to submissive and clinging behavior and fears of separation, beginning by early adulthood and present in a variety of contexts, as indicated by five (or more) of the following:

- (1) has difficulty making everyday decisions without an excessive amount of advice and reassurance from others
- (2) needs others to assume responsibility for most major areas of his or her life
- (3) has difficulty expressing disagreement with others because of fear of loss of support or approval
- (4) has difficulty initiating projects or doing things on his or her own (because of lack of self-confidence in judgment or abilities rather than a lack of motivation or energy)
- (5) goes to excessive lengths to obtain nurturance and support from others, to the point of volunteering to do things that are unpleasant
- (6) feels uncomfortable or helpless when alone because of exaggerated fears of being unable to care for himself or herself
- (7) urgently seeks another relationship as a source of care and support when a close relationship ends
- (8) is unrealistically preoccupied with fears of being left to take care of himself or herself

# Cluster C Personality Disorders

## Dependent Personality Disorder

---

- Prevalence: 2.5-27% of population
  - Sex ratio: F>M
  - Comorbidity: Mood and anxiety disorders, adjustment disorders
  - Family: Cluster C disorders
-

# Cluster C Personality Disorders

## Dependent Personality Disorder

---

- Treatment
    - Psychotherapy - Dynamic, behavior, group, and family therapies are all used successfully
    - Medication - Anxiolytics are often helpful. Antidepressants may be used with comorbid depression.
-

# Cluster C Personality Disorders

## Dependent Personality Disorder

---

- Physician-patient interaction
    - Physicians should take an active role in treatment planning, with clear explanations and recommendations
    - Patients may need encouragement to make decisions about treatment plans
    - Family involvement is often helpful
-

# Cluster C Personality Disorders

## Obsessive Compulsive Personality Disorder

---

- “A pattern of preoccupation with orderliness, perfectionism, and control.” (DSM-IV)
-

# Cluster C Personality Disorders

## Diagnostic Criteria for Obsessive Compulsive Personality Disorder (DSM-IV)

A pervasive pattern of preoccupation with orderliness, perfectionism, and mental and interpersonal control, at the expense of flexibility, openness, and efficiency beginning by early adulthood and present in a variety of contexts, as indicated by four (or more) of the following:

- (1) is preoccupied with details, rules, list, order, organization, or schedules to the extent that the major point of the activity is lost
- (2) shows perfectionism that interferes with task completion (e.g., is unable to complete a project because his or her own overly strict standards are not met)
- (3) is excessively devoted to work and productivity to the exclusion of leisure activities and friendships (not accounted for by obvious economic necessity)
- (4) is overconscientious, scrupulous, and inflexible about matters of morality, ethic, or values (not accounted for by cultural or religious identification)
- (5) is unable to discard worn-out or worthless objects even when they have no sentimental value
- (6) is reluctant to delegate tasks or to work with others unless they submit to exactly his or her way of doing things
- (7) adopts a miserly spending style toward both self and others; money is viewed as something to be hoarded for future catastrophes
- (8) shows rigidity and stubbornness

# Cluster C Personality Disorders

## Obsessive Compulsive Personality Disorder

---

- Prevalence: 1% of population
  - Sex ratio: M:F=2:1
  - Comorbidity: Slight increase in mood and anxiety disorders
  - Family: Obsessive-compulsive personality disorder
-

# Cluster C Personality Disorders

## Obsessive Compulsive Personality Disorder

---

- Treatment
    - Psychotherapy - Psychoanalytic, behavioral, and group therapies are often useful
    - Medication - Serotonin-specific reuptake inhibitor (SSRI) antidepressants may be useful
-



# Cluster C Personality Disorders

## Obsessive Compulsive Personality Disorder

---

- Physician-patient interaction
    - Thorough explanations and specific, detailed information are valued
    - Uncertainty is rarely tolerated
    - Treatment options should be presented with clear risk-benefit analyses
-

# Additional Source Information

for more information see: <http://open.umich.edu/wiki/CitationPolicy>

Slide 4-6: American Psychiatric Association: *Diagnostic and Manual of Mental Disorders*, 4<sup>th</sup> ed, Text Revision (*DSM-IV-TR*), Washington, DC, American Psychiatric Association, 2000, p. 689

Slide 13: Rorschach, NowPublic, <http://www.nowpublic.com/health/roorschach-plate-1-bat-butterfly-moth>

Slide 21: *DSM-IV-TR*, pp. 694

Slide 26: *DSM-IV-TR*, pp. 697

Slide 31: *DSM-IV-TR*, pp. 701

Slide 37: *DSM-IV-TR*, pp. 706

Slide 43: *DSM-IV-TR*, pp. 710

Slide 49: *DSM-IV-TR*, pp. 714

Slide 54: *DSM-IV-TR*, pp. 717

Slide 60: *DSM-IV-TR*, pp. 721

Slide 65: *DSM-IV-TR*, pp. 725

Slide 70: *DSM-IV-TR*, pp. 729