open.michigan

Author(s): Alison Haddock, MD, 2011

License: Unless otherwise noted, this material is made available under the terms of the **Creative Commons Attribution Share Alike 3.0 License**: http://creativecommons.org/licenses/by-sa/3.0/

We have reviewed this material in accordance with U.S. Copyright Law and have tried to maximize your ability to use, share, and adapt it.

Copyright holders of content included in this material should contact **open.michigan@umich.edu** with any questions, corrections, or clarification regarding the use of content.

For more information about **how to cite** these materials visit http://open.umich.edu/privacy-and-terms-use.

Any **medical information** in this material is intended to inform and educate and is **not a tool for self-diagnosis** or a replacement for medical evaluation, advice, diagnosis or treatment by a healthcare professional. Please speak to your physician if you have questions about your medical condition.

Viewer discretion is advised: Some medical content is graphic and may not be suitable for all viewers.





Citation Key

for more information see: http://open.umich.edu/wiki/CitationPolicy

Use + Share + Adapt					
{ Content the copyright holder, author, or law permits you to use, share and adapt. }					
Ø PD-GOV	Public Domain – Government: Works that are produced by the U.S. Government. (17 USC § 105)				
Ø PD-EXP	Public Domain – Expired: Works that are no longer protected due to an expired copyright term.				
Ø PD-SELF	Public Domain – Self Dedicated: Works that a copyright holder has dedicated to the public domain.				
(cc) ZERO	Creative Commons – Zero Waiver				
CC) BY	Creative Commons – Attribution License				
CC BY-SA	Creative Commons – Attribution Share Alike License				
CC BY-NC	Creative Commons – Attribution Noncommercial License				
BY-NC-SA	Creative Commons – Attribution Noncommercial Share Alike License				
③ GNU-FDL	GNU – Free Documentation License				

Make Your Own Assessment

{ Content Open.Michigan believes can be used, shared, and adapted because it is ineligible for copyright. }

PD-INEL Public Domain – Ineligible: Works that are ineligible for copyright protection in the U.S. (17 USC § 102(b)) *laws in your jurisdiction may differ

{ Content Open.Michigan has used under a Fair Use determination. }



Fair Use of works that is determined to be Fair consistent with the U.S. Copyright Act. (17 USC § 107) *laws in your jurisdiction may differ

Our determination **DOES NOT** mean that all uses of this 3rd-party content are Fair Uses and we **DO NOT** guarantee that your use of the content is Fair.

To use this content you should do your own independent analysis to determine whether or not your use will be Fair.

CASE OF THE WEEK

Alison Haddock, PGY4

OBJECTIVES

- Discuss two critical care cases
- Challenges in management and diagnosis
- Review emergent management of a common ED presentation
- Focused exploration of a less common disease process
- Discussion of how the health care system can contribute to individual patient morbidity and mortality

CRITICAL PATIENT IN RESUS BRAVO

- Obese elderly Asian female
- Pale, breathing heavily
- Accompanied by son

ABCS

o A

• Speaking single words

o B

- RR 30
- SpO2 unable to obtain
- C
 - HR 50
 - BP unable to obtain



PULSES AND BPS

• Old ATLS teaching

- SBP > 80 mmHg if palpable radial
- SBP > 70 mmHg if palpable femoral
- SBP > 60 mmHg if palpable carotid
- Not scientifically validated
- Did confirm that loss of pulses occurs in order...
 - radial
 - femoral
 - carotid

Deakin CD and Low JL. Accuracy of the advanced trauma life support guidelines for predicting systolic blood pressure using carotid, femoral, and radial pulses: observational study. BMJ 2000; 321 : 67.

NEXT STEPS

• IV

• Multiple techs attempting

o O2

- Supplemental O2 via NRB
- Monitor
 - Slow HR
 - No BP

NEXT STEPS



(cc) BY-NC-SA robswatski,

robswatski, "Right external jugular vein", flickr

NEXT STEPS

• IV

• 16 gauge in R EJ

o O2

- Supplemental O2 via NRB
- Monitor
 - Slow HR (ranging 40s-50s)
 - No BP (estimate 60-70 SBP)

WHAT NOW?

Brief History

Brief Exam

Further Interventions

More Clinical Data

BRIEF HISTORY

- POD #10 from lap-to-open cholecystecomy
- Prolonged post-operative hospitalization due to "heart problems"
- Discharged home three days ago
- Increasingly weak today
- C/O severe fatigue, "chills"
- Denies measured temps, denies pain
- Taking all medications including coumadin and blood pressure pills



(cc) BY-NC-SA mag3737, flickr

BRIEF EXAM

- Pale, increased WOB
- PERRL, dry MM
- Shallow clear breath sounds
- Slow irregular heartbeat
- Obese/distended and firm abdomen
 - No focal tenderness
- Cool extremities
 - No palpable radial pulse
 - Thready femoral pulse



FURTHER INTERVENTION

- IVF bolus
 - 1L wide open
 - Attempting to obtain additional access
- Pacer pads



WHAT CLINICAL DATA?

- Labs?
- XR?
- CT Scan?
- US?
- Phone-a-friend?





RESUS SHORCUTS

VBG EKG CXR

EKG



PD-INEL Source Unknown

VBG

7.23/44/164

Na 137 K 4.9 Ca 1.04







CXR

DIAGNOSIS?

Shock.

LACTATE & MORTALITY



Mikkelson, M et al. Serum lactate is associated with mortality in severe sepsis independent of organ failure and shock. *Critical Care Medicine*. 2009; 37(5): 1670-1677.

TYPES OF SHOCK

Hypovolemic
Obstructive
Distributive
Cardiogenic

DIFFERENTIAL DIAGNOSIS

- Hypovolemic
 - Hemorrhagic
- Obstructive
 - No apparent evidence of PE, PTX or tamponade...
- Distributive
 - Sepsis from recent hospitalization/surgery
- Cardiogenic
 - Hx of recent cardiac problems
 - Medication toxicity?

- Ongoing bradycardia 40s-50s
- Treatment?
 - For HR <50bpm with evidence of hypoperfusion
 - Or if high risk of progression to complete block
 - Options: atropine vs pacing
- Atropine
 - 0.5 1mg IVP adult dose
 - Anticholinergic positive chronotropic effect
 - Pt has increased HR to 50s-60s without BP improvement

- Respirations increasingly labored
- Abdomen still distended
- Now poorly responsive to son's questioning
- Back to the ABCs!





PhillippN, "Endotracheal tube colored", Wikimedia Commons

- Airway now secure
- Still unable to obtain BP
- Access: single EJ
- Additional 14g placed by tech in Right AC
- Second Liter warmed Normal Saline startedPt started on pressors

PRESSORS (OVER)SIMPLIFIED



• How can we distinguish between types of shock?

RAPID ULTRASOUND IN SHOCK (RUSH)

PUMP
Cardiac
contractility
Camponade
Pneumothorax
RV strain



RAPID ULTRASOUND IN SHOCK (RUSH)

TANK •IVC – size & resp change •FAST



RAPID ULTRASOUND IN SHOCK (RUSH)

PIPES •Aorta **o**DVT





© PD-INEL Source Unknown

ULTRASOUND



Abdomen increasingly distended and firm Surgery contacted

• Treatment initiated for hemorrhagic shock

- O+ pRBCs placed on rapid transfuser
 - Massive transfusion anticipated
 - Given calcium chloride

REVERSAL OF ANTICOAGULATION: FIRST STEPS

• FFP

- Typical adult pt requires 3-4 units to reverse
- Contains all vitamin K dependent factors
- Does not fully reverse
 - Ex: factor IX does not rise >20% of normal post FFP (not reflected in INR)
- Requires thawing
- Risks volume overload
- o Vitamin K
 - 10mg slow IV
 - Starts to work in 4hrs

REVERSAL OF ANTICOAGULATION: NEXT STEPS

• Recombinant Activated Factor VII

- Expensive, limited literature
- UofM: "serious bleed associated with prolonged INR after significant clotting factor replacement"
- 1200mcg x one dose
- Prothrombin Complex Concentrate
 - Plasma-derived product, no matching required
 - Virally inactivated and 20x less volume than FFP
 - Contains II, IX, and X (+ VII in UK)
 - Currently infrequently used in US
 - Expensive
 - Potentially thrombogenic

Rapid transfusion of 3U FFP, 4U pRBCs Surgery at bedside

• Still no BP on max dose dopamine (20mcg/kg/min)

• Pulse check = no carotid or femoral

• PEA

- Compressions initiated
- Single dose of 1mg epinephrine
- Return of strong pulse

• ABCs

- Airway secured with ETT
- Ongoing hypotension despite pressors
- o Volume
 - Cordis inserted into R groin + 14G PIV + 16G PIV
 - Femoral arterial line
 - Rapid infuser for pRBCs, FFP
- Pressors
 - Max dose dopamine
 - Epinephrine and norepinephrine initiated post-arrest

LABS

Hematology:			
CBCP:			
10.1	Cardiac Enzymes (18:00):		
26.2 >< 375	Troponin: 0.11; CPK: 102; CKMB: 2.3;		
31.3			
	Liver Panel:		
Diff (Automatic):	Prot: 5.7; Alb: 3.1;		
NEUT LYMPH MONO EOS BASO	AST: 54; ALT: 23;		
%:83.7 8.3 8.0 .0 .0	TBILI: 1.7;		
#:21.9 2.2 2.1 0.0 0.0	ALK: 156;		
Chemistry:			
	Coags:		
142c 102 24 Cal: 9.32c AnGap: 17	INR: 1.8 (PT: 18.1); PTT: 24.8;		
< 190 BUN/Cr: 16			
4.6 22 1.5 CalcOsm: 301			

- Mismatch between BPs
 - Femoral arterial line = SBPs in 60s
 - Cuff pressure = SBPs in 100s
- Overall poor responsiveness to pressors and fluids
- Additional diagnosis made:





Derek K. Miller, flickr

• Definition

- Sustained intraabdominal pressure of >20 mmHg associated with new organ dysfunction or failure
- Measurement of Intra-Abdominal Pressure
 - Challenging clinical diagnosis
 - Direct peritoneal cannulation, rectal, gastric, IVC
 - Most popular = bladder
 - Routinely tracked in ICU settings
- Relatively high incidence
 - One study found ACS in 14% of high-risk trauma pts
 - Another found 50% of ICU pts had IAH (>12 mmHg)

- Primary ACS: injury/disease in abdomen
 - Abdominal trauma
 - Abdominal hemorrhage
 - Bowel obstruction
 - Intraperitoneal sepsis
 - Ruptured AAA
 - Acute pancreatitis
 - Less acute: morbid obesity, pregnancy, massive ascites
- Secondary ACS: third-spacing in abdomen
 - Severe sepsis, burns
 - Any shock requiring massive fluid resuscitation

Box 1: Emergency department patients at risk of developing intra-abdominal hypertension (IAH) and abdominal compartment syndrome (ACS)

Patients who should be considered at higher risk of developing IAH and ACS:^{35 36}

- Patients with open or blunt abdominal trauma
- Patients requiring large volume fluid resuscitation (especially in the context of an underlying capillary leak problem)—for example:
 - pancreatitis
 - septic shock
 - trauma
 - severe burns
- Patients with increased intra-luminal contents
 - gastroparesis
 - ileus

Ø PD-INEL

- Patients with increased abdominal contents
 - haemoperitoneum or pneumoperitoneum
 - ascites or liver dysfunction

Harrison, SE, et al. Abdominal compartment syndrome: an. emergency department perspective. *Emerg Med J* 2008;25:128–132

	Increased	Decreased	No change
Mean blood pressure	-	-	×
Heart rate	×	-	-
Peak airway pressure	×	-	-
Thoracic/pleural pressure	×	-	-
Central venous pressure	×	-	-
Pulmonary capillary wedge pressure	×	-	-
Inferior vena cava pressure	×	-	-
Renal vein pressure	×	-	-
Systemic vascular resistance	×	-	-
Cardiac output	-	×	-
Venous return	-	×	-
Visceral blood flow	-	×	-
Gastric mucosal pH		×	
Renal blood flow	-	×	-
Glomerular filtration rate	-	×	-
Cerebrospinal fluid pressure	×		
Abdominal wall compliance	-	×	-

Table 40.1. Physiological consequences of intra-abdominal hypertension

SCHEIN'S COMMON SENSE EMERGENCY ABDOMINAL SURGERY 2009, Part 2, 435-443

Ø PD-INEL



Fig. 40.1. The abdominal compartment syndrome

SCHEIN'S COMMON SENSE EMERGENCY ABDOMINAL SURGERY

2009, Part 2, 435-443

• Definitive management: surgical decompression

Box 2: Conservative measures for treatment of abdominal compartment syndrome

- E Paracentesis⁵⁹
- Gastric and rectal suctioning
- Prokinetic agents (metoclopramide, domperidone, erythromycin, prostygmine)
- Diuretic therapy
- Continuous veno-venous haemofiltration with aggressive ultrafiltration
- ► Sedation and paralysis⁶⁰
- Body positioning

Harrison, SE, et al. Abdominal compartment syndrome: an. emergency department perspective. *Emerg Med J* 2008;25:128–132

Ø PD-INEL

• Definitive management: surgical decompression

• In ED, pt difficult to ventilate

- Given doses of versed and vecuronium
- Anesthesia arrived to assist
- Covered with piperacillin/tazobactamOG Tube placed before transport to OR

OPERATIVE REPORT

INDICATIONS FOR THE PROCEDURE: Mrs. PG is an elderly woman who underwent a laparoscopic converted to open cholecystectomy 10 days ago. She presented to the ER with complaints of feeling weak and light-headed. Shortly after admission to the ER, she became hypotensive and had a PEA arrest. She was rescuscitated and a FAST scan showed a large amount of fluid in her abdomen; her abdomen was also quite tense on examination. She is anti-coagulated with coumadin and her INR at that time was 1.8. She was presumed to have an intra-abdominal bleed and was therefore taken emergently to the OR. She was requiring significant amounts of fluid rescuscitation, including 4 units of PRBC's as well as pharmacologic pressure support. She was intubated in the ER.

PRCEDURE: Time out was performed, confirming correct patient. Anesthesia was induced with general endotracheal anesthesia. The patient was positioned in the supine position on the table and was prepped and draped in the usual aseptic fashion. Her abdomen was then entered through a midline incision extending from xiphoid to just above the pubis. Dissection was carried own through the subcutaneous tissues and the fascia was opend in the provine. Upon entering the peritoneal cavity, a large amount of blood and clot y encountered and evacuated. Approximately 1.500 cc's of clot and old blood was evacuated, the majority of the clot was encountered inferior to the liver. This was evacuated and there was obvious bleeding from the gallbladder fossa. This was controlled with a combination of electrocautery, fibrillar, and surgicel. The area was then widely irrigated with warm normal saline and topical thrombin spray was applied and packs were then placed. Five minutes elapsed and we re-examined the area and no further bleeding was appreciated. At this point, all packs were removed. There was no further evidence of bleeding. Since the patient had undergone massive fluid rescusitation and anesthesia was already having some difficulties with ventilation, we elected to leave her abdomen open and place and abdominal VAC. Sponge and needle counts were correct at the conclusion of the case. The patient was taken to the SICU in critical condition.

Upon entering the peritoneal cavity, a large amount of blood and clot was encountered and evacuated. Approximately 1,500 cc's of clot and old blood was evacuated, the majority of the clot was encountered inferior to the liver. This was evacuated and there was obvious bleeding from the gallbladder fossa.

we elected to leave her abdomen open and place an abdominal VAC

HOSPITAL COURSE

- Extensive ICU course
 - ARF with anuria, on CRRT
 - Resp failure requiring tracheostomy
 - Febrile with +BAL (stenotrophomonas), MDR UTI, infected rectus sheath hematoma (both E coli)
 - Intermittent A-fib with RVR
 - Multiple pulmonary emboli
- Transferred SICU to BICU to CCMU
- Discharged home two months later with PEA as primary diagnosis

52yo M with a hx of "liver and kidney problems"CC: SOB

- Arrives in Resus A in acute distress
- Gasping for breath, saying single words
- Pale, diaphoretic
- Bradycardic with palpable radial pulse
- Unable to measure BP or SpO2
- PIV placed by EDT

• Becomes unresponsive <60sec after arrival

- BVM applied
- Given atropine 1mg IV w/o change in status
- Loses pulses
- CPR initiated
- 2 rounds epinephrine/atropine
- Intubated w/o meds

- Empirically medicated given "kidney" history
 - Calcium gluconate
 - Sodium bicarbonate
- VBG returns with K of 8.0
 - Started insulin and glucose
 - Albuterol via ETT
- ROSC after <10 minutes (3 rounds)
- Spontaneous movements observed

- Gradual onset of hypotension after ROSC
- Started on dopamine
- Empiric sepsis coverage
 - Piperacillin/tazobactam
 - Vancomycin
- Accepted for admission by CCMU
- Sedated with propofol

• Cooling?

- 2 week admission to CCMU requiring dialysis
- Normal neurologic status post-extubation
- Discharged home with outpt dialysis, otherwise doing well

THANK YOU!



(CC) BY-NC-SA

zoomar, "Emergency Sign, Ballard Swedish Hospital", flickr,CC: BY-NC-SA 2.0,

Additional Source Information

for more information see: http://open.umich.edu/wiki/CitationPolicy

Slide 7, Image 1: OldakQuill, "Pioneer plaque line-drawing of a human male", Wikimedia commons, http://commons.wikimedia.org/wiki/ File:Pioneer_plaque_line-drawing_of_a_human_male.svg, Public Domain - Government

Slide 10, Image 1: robswatski, "Right external jugular vein", flickr, http://www.flickr.com/photos/rswatski/5913031286/, CC: BY-NC-SA 2.0, http:// creativecommons.org/licenses/by-nc-sa/2.0/deed.en.

Slide 13-16, Image 1: mag3737, "Emergency", flickr, http://www.flickr.com/photos/mag3737/3572764826/, CC: BY-NC-SA 2.0, http:// creativecommons.org/licenses/by-nc-sa/2.0/deed.en.

Slide 18, Image 1: Source Unknown.

Slide 20, Image 1: Source Unknown.

Slide 22, Image 1: Mikkelson, M et al. Serum lactate is associated with mortality in severe sepsis independent of organ

failure and shock. Critical Care Medicine. 2009; 37(5): 1670-1677.

Slide 26, Image 1: PhillippN, "Endotracheal tube colored", Wikimedia Commons, http://commons.wikimedia.org/wiki/ File:Endotracheal_tube_colored.png, Public Domain - Government.

Slide 30, Image 1: Source Unknown.

Slide 31, Image 1: Source Unknown.

Slide 32, Image 1: Source Unknown.

Slide 33, Image 1: Source Unknown.

Slide 41, Image 2: Derek K. Miller, "Brined turkey experiment 3", flickr, http://www.flickr.com/photos/penmachine/4185510308/, CC: BY-NC 2.0, http://creativecommons.org/licenses/by-nc/2.0/deed.en.

Slide 44, 48, Table 1: Harrison, SE, et al. Abdominal compartment syndrome: an. emergency department perspective.

Emerg Med J 2008;25:128-132

Slide 45, 46, Image 1: SCHEIN'S COMMON SENSE EMERGENCY ABDOMINAL SURGERY, 2009, Part 2, 435-443

Additional Source Information

for more information see: http://open.umich.edu/wiki/CitationPolicy

Slide 57, Image 2: zoomar, "Emergency Sign, Ballard Swedish Hospital", flickr, http://www.flickr.com/photos/zoomar/468637568/, CC: BY-NC-SA 2.0, http://creativecommons.org/licenses/by-nc-sa/2.0/deed.en